



Household water treatment

Household-based approaches to water treatment may be more effective and cost-effective means of preventing diarrhoeal disease than conventional treatment at source. This fact sheet summarizes the available evidence and some of the leading approaches.

Diarrhoeal diseases kill an estimated 2.5 million people each year, the majority children under five. They are also the third highest cause of morbidity. As a result, health authorities generally accept that safe water plays an important role in preventing outbreaks of diarrhoeal disease.

Water quality: refining the dominant paradigm

Fifteen years ago, Esrey and colleagues reviewed previous studies on the impact of environmental interventions on diarrhoea, and found improvements in water quality to be considerably less effective than those aimed at water quantity, water availability, sanitation, and hygiene interventions (Esrey et al., 1991). Ubiquitously cited in both professional journals and practical guides, the reviews have led to the dominant paradigm respecting water supply and sanitation interventions: that to achieve broad health impact, greater attention should be given to safe excreta disposal and proper use of water for personal and domestic hygiene rather than to drinking-water quality.

Recently, however, an increasing body of evidence has suggested the need to refine the dominant paradigm (Clasen & Cairncross, 2004). Much of this evidence arises from a relatively new approach to enhancing water quality as part of a public health initiative: improved household water management and storage. Esrey's conclusions regarding the impact of water quality on diarrhoeal disease were based exclusively on studies involving interventions at the point of distribution. However, it is well known that even safe water becomes faecally contaminated during collection, transport, storage and drawing in the home (Wright et al., 2004). Accordingly, improving water quality at the point of distribution only may not secure the full health gains that are possible by ensuring that drinking water is microbiologically safe at the point of use (POU). This distinction was suggested in a recent systematic review that demonstrated a 34% reduction in diarrhoea from interventions at the point of use, fully twice the impact reported by Esrey of improvements at the source (Fewtrell et al., 2005).

The health impact of treating water at the point of consumption, however, is not universal or absolute. Except in the case of *Vibrio cholerae*, a reduction in waterborne pathogens is not clearly associated with a corresponding reduction in diarrhoea (Gundry et al., 2004). A pending Cochrane review should clarify the apparent difference in health impact between interventions at point of distribution versus those at the point of use, and help explain the heterogeneous results observed for interventions at the household level.

Household water treatment and the WHO

As part of its Millennium Development Goals, the United Nations expressed its commitment by 2015 to reduce by one half the 1.1 billion people without sustainable access to improved water supply. Providing safe, piped-in, disinfected water, to each household may be the best solution to waterborne disease. The WHO acknowledges however, that such a solution would entail an



Household water treatment

investment of tens of billions of dollars each year. Accordingly, it has called for other approaches while progress is made in improving infrastructure.

According to the 2002 World Health Report, point-of-use water treatment, such as household-based chlorination, is the most cost-effective environmental intervention to prevent diarrhoeal disease across a wide range of countries and settings (WHO, 2002). The WHO also commissioned a comprehensive review to identify the most promising POU treatment technologies based on selected technical characteristics and performance criteria, including effectiveness in improving and maintaining microbial water quality, health impact, technical difficulty or simplicity, accessibility, cost, acceptability, sustainability and potential for dissemination (Sobsey, 2002). After evaluating at least 37 different technologies, Sobsey concluded that five were the most promising: filtration with ceramic filters, chlorination with storage in an improved vessel, solar disinfection in clear bottles, thermal disinfection (pasteurization) in solar cookers or reflectors, and combination systems employing chemical flocculation and chlorination.

While this fact sheet focuses on these technology groups, readers are urged to explore other options which may be more suitable for a particular setting. Moreover, the potential commercial market for household-based water treatment has attracted private sector participants who are attempting to adapt or develop new technologies. Accordingly, readers are encouraged to investigate these emerging technologies.

Certain technologies

Chemical disinfection

Chemical disinfection is the most widely-practised means of treating water at the community level. It is also the method used most broadly in the home. While a wide range of oxidants are used in treating water, most household-based interventions employ free chlorine derived from liquid sodium hypochlorite or solid calcium hypochlorite which are usually available and affordable. Tablets formed from chlorinated isocyanurates (e.g., NaDCC), a leading emergency treatment of drinking water, and novel systems for on-site generation of oxidants such as chlorine dioxide, may also have a role in household water treatment in the future. At doses of a few mg/l and contact time of about 30 minutes, free chlorine inactivates more than 4 logs of enteric pathogens, the notable exceptions being *Cryptosporidium* and *Mycobacterium* species. The "Safe Water System", a programmatic intervention developed by the US Centers for Disease Control and Prevention that combines chlorination of water in the home with safe storage and hygiene instruction, has an estimated 5 million users in 19 countries (www.cdc.gov/safewater/default.htm). Its impact in reducing diarrhoeal diseases has been documented (Quick et al., 2002). Like most other household-based water interventions, however, the hardware must be accompanied by an extensive behavioural change programme to stimulate adoption and continued utilization by householders.

Filtration

Household filters potentially present certain advantages over other technologies. They operate under a variety of conditions (temperature, pH, turbidity), introduce no chemicals into the water that may affect use due to objections about taste and odour, are easy to use, and improve the water aesthetically, thus potentially encouraging routine use without extensive intervention to promote behavioural change. Higher quality ceramic filters treated with bacteriostatic silver have been shown effective in the lab at reducing waterborne protozoa by more than 3 logs and bacteria by more than 6 logs. Their potential usefulness as a public health intervention has been suggested in a recent field trial (Clasen et al., 2004). While the up-front cost of gravity systems employing such commercial ceramics is high (US\$10 to US\$25), their long life (up to 50,000 litres per ceramic candle element) renders such systems comparable to chlorination on a per litre treated basis. The improving quality of locally-fabricated silver coated ceramics is particularly promising as a sustainable and low-cost alternative. Slow-sand filters, which remove suspended solids and microbes by means of a slime layer (*schmutzdecke*) that develops within the top few centimetres of sand, are capable of removing 2 logs or more of enteric pathogens, if properly constructed, operated and maintained (Hijnen et al, 2004). A simpler but more advanced version, known as the bio-sand filter, was specifically designed for intermittent use and is more suitable for household applications. It has been tested (Palmeteer et al., 1999) and is being deployed widely in development settings with the help of CAWST, a Canadian NGO. Other filtration media, ranging from simple folded sari cloth for the removal of cholera-associated zooplankton

(Colwell et al., 2003) to advanced but inexpensive carbon nanofibre membranes capable of removing even viruses at gravity pressure, demonstrate the broad range of opportunities for filtering water at the household level.

Thermal and solar disinfection

Boiling or heat treatment of water with fuel is effective against the full range of microbial pathogens and can be employed regardless of the turbidity or dissolved constituents of water. While the WHO and others recommend bringing water to a rolling boil, this is mainly intended as a visual indication that a high temperature has been achieved. In fact, studies have demonstrated that heating to pasteurization temperatures (60°C) for 10 minutes will kill or deactivate most pathogens. However, the cost and time used in procuring fuel, and the environmental issues around denuding forests and adding to poor indoor air quality, have led to other alternatives. Solar disinfection, which combines thermal and UV radiation, has repeatedly shown to be effective for eliminating microbial pathogens (Reed, 2004) and reduce diarrhoeal morbidity (Conroy et al., 1999). Among the most practical and economical is the 'Sodis' system, developed and promoted by the Swiss Federal Institute for Environmental Science and Technology (EAWAG). It consists of placing low turbidity (<30NTU) water in clear plastic bottles (normally discarded 2 litre beverage bottles, preferably PET) after aerating it to increase oxygenation and exposing the bottles to the sun, usually by placing them on corrugated metal roofs. Exposure times vary from 6 to 48 hours depending on the intensity of sunlight. Like filters, thermal and solar disinfection do not provide residual protection against recontamination. Accordingly, householders must have a sufficient number of bottles to allow them to cool and maintain treated water in the bottles until it is actually consumed.

Combination flocculation and disinfection

A challenge for most household-based water treatment technologies is high turbidity. Solids can use up free chlorine and other chemical disinfectants, cause premature clogging of filters, and block UV radiation essential in solar disinfection. While turbidity can often be managed by pre-treatment or even simple sedimentation, flocculation/coagulation using additives such as alum can be an effective and relatively low-cost option. Such forms of assisted sedimentation have been shown to reduce the levels of certain microbial pathogens, especially protozoa which may otherwise present a challenge to chemical disinfectants. However, disinfection is still required in most cases for complete microbial protection. Certain manufacturers have combined flocculation and time-released disinfection in a single product that is sold in sachets for household use. Proctor & Gamble's PUR® product, the most extensively tested, has been shown to reduce waterborne cysts by more than 3 logs, viruses by more than 4 logs and bacteria by more than 7 logs. Unlike the other methods of household water treatment, it has also been shown to be effective in reducing arsenic, an important non-microbial contaminant in certain settings, by more than 2 logs. Field studies have demonstrated such flocculation-disinfection products effective in preventing diarrhoeal diseases (Reller et al., 2003). While these products are relatively expensive on a per-litre treated basis, they may have application in certain emergency and other settings. It may also be possible to achieve similar results by combining conventional and lower cost approaches to assisted sedimentation and subsequent disinfection at the expense of convenience.

Acceptability, affordability and sustainability

Household water treatment as an intervention against diarrhoeal disease is still at a nascent stage in its development. Questions about acceptability, affordability, long-term utilization and sustainability must still be addressed, particularly in programmatic settings. These issues will ultimately help determine the potential role of household water treatment in preventing diarrhoeal disease among vulnerable populations.

References and further reading

- Clasen, T, Cairncross S (2004). Household water treatment: refining the dominant paradigm. *J Trop Med Hyg* 9:187–191.
- Clasen T, Brown J, Suntura O, Collin S (2004). Reducing diarrhoea through household-based ceramic filtration of drinking water: a randomized, controlled trial in Bolivia. *Am J Trop Med Hyg* 70(6):651–7.
- Colwell R R, Huq A, Islam M S, Aziz K M A, Yunus M, Khan N H, Mahmud A, Sack R B, Nair G B, Chakraborty J, Sack D A, Russek-Cohen E (2003). Reduction of cholera in Bangladeshi villages by simple filtration. *Proc Nat Acad Sci* 100(3): 1051–5.
- Conroy R M, Meegan M E, Joyce T, McGuigan K, Barnes J (1999). Solar disinfection of water reduces diarrhoeal disease: an update. *Arch Dis Child* 81(4): 337–8.
- Esrey S A, Potash J B, Roberts L & Schiff, C (1991). Effects of improved water supply and sanitation on ascariasis, diarrhoea, dracunculiasis, hookworm infection, schistosomiasis, and trachoma. *Bull WHO* 69, 609–21.

Household water treatment

- Fewtrell L, Kaufmann R, Kay D, Enanoria W, Haller L, Colford J (2005). Water, sanitation, and hygiene interventions to reduce diarrhoea in less developed countries: a systematic review and meta-analysis. *Lancet Infect Dis* 5:42–52.
- Gundry S, Wright J, Conroy R (2004). A systematic review of the health outcomes related to household water quality in developing countries. *J Water & Health* 2(1):1–13 In press.
- Hijnen W A, Schijven J F, Bonne P, Visser A, Medema G J (2004). Elimination of viruses, bacteria and protozoan oocysts by slow sand filtration. *Water Sci Technol* 50(1):147–54.
- Palmateer, G., Manz, D, Jurkovic, A, McInnis, R, Unger, S, Kwan, K K, Dudka, B J (1999). Toxicant and parasite challenge of Manz intermittent slow sand filter. *Environ Toxicol* 14:217–225.
- Quick R E, Kimura A, Thevos A, Tembo M, Shamputa I, Hutwagner L, Mintz E (2002). Diarrhea prevention through household-level water disinfection and safe storage in Zambia. *Am J Trop Med Hyg* 66(5): 584–9.
- Reller M E, Mendoza C E, Lopez M B, Alvarez M, Hoekstra R M, Olson C A, Baier K G, Keswick B H, Luby S P (2002). A randomized controlled trial of household-based flocculant-disinfectant drinking water treatment for diarrhea prevention in rural Guatemala. *Am J Trop Med Hyg* 64(4).
- Reed R H (2004). The inactivation of microbes by sunlight: solar disinfection as a water treatment process. *Adv Appl Microbiol* 54:333–65.
- Sobsey MD (2002). Managing water in the home: accelerated health gains from improved water supply. Geneva: *The World Health Organization* (WHO/SDE/WSH/02.07).
- WHO (2002). The World Health Report 2002. Geneva: The World Health Organization.
- Wright J, Gundry S, Conroy (2003). Household drinking water in developing countries: a systematic review of microbiological contamination between source and point-of-use. *Trop Med & Int'l Health* 9(1):106–17.

Author: T Clasen

Quality assurance: S Cairncross

Design: S Plummer

Illustrations: R Shaw

WELL fact sheets are a series of short documents containing recommendations made by WELL, written primarily as a response to WELL enquiries and frequently-asked questions. These are regularly updated to reflect current issues and thinking within the water, sanitation and environmental health sector.

WELL Network Partners are developing regional annexes to a number of existing fact sheets. For details, refer to the WELL website: <http://www.lboro.ac.uk/well/about-well/about-well.htm>

WELL is a resource centre network providing access to information and support in water, sanitation and environmental health for the Department for International Development (DFID) of the British Government.

For further details, contact:

WELL at WEDC, Loughborough University, Leicestershire, LE11 3TU, UK.

Phone: +44 1509 222885 Fax: +44 1509 211079 E-mail: well@lboro.ac.uk

<http://www.lboro.ac.uk/well/>